

University of Groningen

Dilated Virchow-Robin spaces in primary open-angle glaucoma

Wostyn, Peter; De Groot, Veva; Van Dam, Debby; Audenaert, Kurt; Killer, Hanspeter Esriel;
De Deyn, Peter Paul

Published in:
Acta radiologica open

DOI:
[10.1177/2058460116653630](https://doi.org/10.1177/2058460116653630)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2016

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Wostyn, P., De Groot, V., Van Dam, D., Audenaert, K., Killer, H. E., & De Deyn, P. P. (2016). Dilated Virchow-Robin spaces in primary open-angle glaucoma: a biomarker of glymphatic waste clearance dysfunction? *Acta radiologica open*, 5(8). <https://doi.org/10.1177/2058460116653630>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Dilated Virchow-Robin spaces in primary open-angle glaucoma: a biomarker of glymphatic waste clearance dysfunction?

Peter Wostyn¹, Veva De Groot², Debby Van Dam^{3,4},
Kurt Audenaert⁵, Hanspeter Esriel Killer⁶
and Peter Paul De Deyn^{3,4,7}

Acta Radiologica Open
5(8) 1–2
© The Foundation Acta Radiologica
2016
Reprints and permissions:
sagepub.co.uk/journalsPermissions.nav
DOI: 10.1177/2058460116653630
arr.sagepub.com

We read with great interest the article entitled “Ocular blood flow and cerebrospinal fluid pressure in glaucoma” by Promelle et al. (1). The authors review the most recent research on alterations in ocular blood flow and/or cerebrospinal fluid (CSF) flow in glaucoma. They state that blood–CSF flow interactions may be involved in glaucoma, and also note that “the CSF’s behavior is supposedly linked to blood pulsatility.” We fully agree with this notion, as we believe that the vascular system could play a role in the pathogenesis of glaucoma through a mechanical interplay with the CSF circulation.

The subarachnoid space (SAS) of the optic nerve (ON) is contiguous with the SAS of the brain in a normal population. Growing evidence in the literature provides strong support for the concept that CSF pressure and composition in the SAS surrounding the ON may have fundamental significance in the pathogenesis of glaucoma (2,3). As an extension of the brain, the ON displays remarkable similarities to the brain in terms of anatomy and functionality (4). Knowledge obtained from brain research could therefore lead to new insights into the ON and vice versa. Recently, the “glymphatic system” of the brain has been discovered in rodents by Iliff et al. (5). The authors argued that this system was critical to the efficient clearance of interstitial solutes, including amyloid- β , from the brain (5). Their findings suggested a brain-wide paravascular pathway in which CSF flows from the SAS along the arteries and arterioles into the paravascular Virchow-Robin spaces (VRSs) to exchange with interstitial fluid (ISF), and ISF is cleared from the brain into the VRSs surrounding the exiting veins. As ISF exits the brain through the paravenous route, it travels to the lymphatic vessels in the neck, and eventually returns its contents to the systemic circulation.

Paravascular CSF recirculation and ISF solute clearance is dependent upon water transport via astroglial aquaporin-4 water channels, which are localized to

perivascular astrocytic endfeet ensheathing the cerebral vasculature (5). Importantly, there is also evidence demonstrating that there is a mechanical interplay between the intracranial vascular system and the glymphatic system. Indeed, a recent study demonstrated that cerebral arterial pulsatility is a key driver of paravascular CSF influx and subsequent CSF-ISF exchange in the brain (6). Loss of elasticity in aging cerebral arteries with progressive atherosclerosis, or cessation of pulsations, might contribute to failure of the clearance of interstitial waste, including amyloid- β , from the brain, and might play a role in the pathogenesis of Alzheimer’s disease (6).

The above findings indicate that the VRSs have an important role in the homeostasis of cerebral fluids in the central nervous system (7), and suggest a possible correlation between VRS enlargement and a disturbance of CSF dynamics. Indeed, if CSF outflow is reduced as a consequence of either CSF flow obstruction or cerebral artery pulsatility inefficiency, or cerebrospinal venous insufficiency and lymphatic disorders, local perivascular CSF recirculation may be impaired and, consequently,

¹Department of Psychiatry, PC Sint-Amandus, Beernem, Belgium

²Department of Ophthalmology, Antwerp University Hospital, Antwerp, Belgium

³Laboratory of Neurochemistry and Behavior, Institute Born-Bunge, University of Antwerp, Department of Biomedical Sciences, Antwerp, Belgium

⁴Department of Neurology and Alzheimer Research Center, University of Groningen and University Medical Center Groningen, Groningen, The Netherlands

⁵Department of Psychiatry, Ghent University Hospital, Ghent, Belgium

⁶Department of Ophthalmology, Kantonsspital Aarau, Aarau, Switzerland

⁷Department of Neurology and Memory Clinic, Middelheim General Hospital (ZNA), Antwerp, Belgium

Corresponding author:

Peter Wostyn, Department of Psychiatry, PC Sint-Amandus, Reigerlostraat 10, 8730 Beernem, Belgium.
Email: wostyn.peter@skynet.be



the VRSs may dilate due to fluid retention (7). Interestingly, dilated VRSs are a known histological feature of cerebral small vessel disease and can be demonstrated on magnetic resonance imaging (8).

Since the ON is a direct extension of the brain, the question is whether there is also evidence for the existence of a paravascular transport system in the ON. The observation of such an anatomically distinct clearing system in the ON could also provide new insights into the pathogenesis of glaucoma. Indeed, if confirmed, one might expect that a dysfunctional glymphatic system could ultimately result in reduced neurotoxin clearance in the ON and lead to glaucomatous neurodegeneration (4).

Many studies have demonstrated vascular-related risk factors for glaucoma. Atherosclerotic cerebrovascular disease is a risk factor for glaucomatous optic neuropathy, and both systemic hypertension and systemic arterial stiffness have been reported to be associated with primary open-angle glaucoma (POAG) (9,10). Given that impairment of the arterial pulsation-driven “perivascular pump” may lead to CSF flux disturbance, and given that distension of the VRSs may be related to the fluid retention in and along the paravascular circulation (7), we postulate that dilated VRSs could occur in POAG in association with arterial stiffening or loss of arterial pulsatility. Interestingly, in accordance with this view, a very recent study showed significantly higher deep white matter lesion load and increased numbers of dilated VRSs in association with POAG suggesting the presence of significant cerebral small vessel disease (8). If dilation of VRSs may result from a disturbed interplay between the vascular and glymphatic system, then preventing arterial stiffness may protect against glaucoma, at least in part, by promoting solute clearance via the CSF.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

1. Promelle V, Daouk J, Bouzerar R, et al. Ocular blood flow and cerebrospinal fluid pressure in glaucoma. *Acta Radiol Open* 2016;5:2058460115624275.
2. Berdahl JP, Allingham RR. Intracranial pressure and glaucoma. *Curr Opin Ophthalmol* 2010;21:106–111.
3. Killer HE, Miller NR, Flammer J, et al. Cerebrospinal fluid exchange in the optic nerve in normal-tension glaucoma. *Br J Ophthalmol* 2012;96:544–548.
4. Wostyn P, Van Dam D, Audenaert K, et al. A new glaucoma hypothesis: a role of glymphatic system dysfunction. *Fluids Barriers CNS* 2015;12:16.
5. Iliff JJ, Wang M, Liao Y, et al. A paravascular pathway facilitates CSF flow through the brain parenchyma and the clearance of interstitial solutes, including amyloid β . *Sci Transl Med* 2012;4:147ra111.
6. Iliff JJ, Wang M, Zeppenfeld DM, et al. Cerebral arterial pulsation drives paravascular CSF-interstitial fluid exchange in the murine brain. *J Neurosci* 2013;33:18190–18199.
7. Gallina P, Scollato A, Conti R, et al. A β Clearance, “hub” of multiple deficiencies leading to Alzheimer disease. *Front Aging Neurosci* 2015;7:200.
8. Mercieca K, Cain J, Hansen T, et al. Primary open angle glaucoma is associated with MR biomarkers of cerebral small vessel disease. *Sci Rep* 2016;6:22160.
9. Nakazawa T. Ocular blood flow and influencing factors for glaucoma. *Asia Pac J Ophthalmol (Phila)* 2016;5:38–44.
10. Shim SH, Kim CY, Kim JM, et al. The role of systemic arterial stiffness in open-angle glaucoma with diabetes mellitus. *Biomed Res Int* 2015;2015:425835.